


Kronofer[®] INNOVATIVE THREE-LAYER IRON TABLETS WITH VITAMIN C

Prevention and treatment of iron deficiency anaemia, increased iron requirement.

WHAT	The solution for an effective iron supplementation
WHEN	To raise the physiological levels of iron.
WHY	To treat iron deficiency and regain vitality.





DOSAGE 1 tablet /day preferably in the morning.

COMPOSITION PER TABLET

"FAST" layer - release within 5 minutes

	mg/tablet
Iron bisglycinate	10

"NORMAL" layer - release within 45 minutes

	mg/tablet
Iron sulphate	10
Vitamin C	70

"RETARD" layer - release within 8 hours

	mg/tablet
Iron pyrophosphate	10



IRON SUPPLEMENTATION

Kr⊘nofer[®]

INNOVATIVE THREE-LAYER IRON TABLETS WITH VITAMIN C

Three bioavailable sources of iron targeted at the best absorption site.

Innovative technology: Time and Site Controlled release.



3 PHASES tablet release



FERROUS BISGLYCINATE, IRON-CHELATED AMINO ACIDS¹ FAST layer: Release in 5 minutes – release during the first tract of duodenum for maximum gastric tolerability, prompt release and quick absorption.



● FERROUS SULPHATE, A BIOAVAILABLE SOURCE OF IRON² NORMAL layer: Release in 45 minutes – release during the duodenal transit bypassing the gastric release, avoiding the most common problems on intolerance to iron sulphate.

VITAMIN C

To help increase iron absorption and strengthen the immune system.

● FERRIC PYROPHOSPHATE, pH INDEPENDENT, MICROENCAPSULATED IRON³ **RETARD layer:** Release in 8 hours – release during the jejunal transit to increase absorption in the latter part of intestine.

1- Piñeda O. et al.: Effectiveness of iron amino acid chelate on the treatment of iron deficiency anemia in adolescents. J Appl Nutr 1994; 46:2-13. 2- Christides T. et al.: Iron bioavailability from commercially available iron supplements. Eur J Nutr. 2015; 54(8):1345-52. 3- Iron statis improvement in LIPOFER fortified juice vs.placebo in woman British journal of nutrition 2011, 105 1652-1659

ENSURING BETTER PATIENT COMPLIANCE:

• High absorption

Less gastric irritation

- High bioavailability
 Absence of unpleasant

metallic taste



Manufactured by: SIIT srl Via Ariosto 50/60 20090 Trezzano sul Naviglio - Milan Italy Tel· +39 02 48 43 52 1 Email: info@siit-it.com

Distributed by: PHARMACON

PHARMACON (PTE) LTD 8 Kaki Bukit Road 2 #04-27/28 Ruby Warehouse Complex Singapore 417841 Tel: +65 6741 8500 Email: enquiry@pharmacon.com.sg